

Submitted: 27/09/2014

Accepted: 24/01/2015

Published: 08/02/2015

## Review on mechanisms of dairy summer infertility and implications for hormonal intervention

B.U. Wakayo<sup>1,\*</sup>, P.S. Brar<sup>2</sup> and S. Prabhakar<sup>2</sup>

<sup>1</sup>*College of Veterinary Medicine-Jigjiga University, P.O.Box 1020, Jigjiga Ethiopian Somali Regional State, Ethiopia*

<sup>2</sup>*Department of Veterinary Gynaecology and Obstetrics, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana-141 004, India*

### Abstract

In dairy cows and buffaloes, summer heat stress (HS) reduces milk yield and delays return to pregnancy leading to financial loss. Clues for effective interventions against summer infertility (SI) lie in understanding the underlying mechanisms. This article reviews current knowledge on the mechanisms of bovine SI and their implication for hormonal management. Under HS dairy animals encounter anestrus, silent cycles and repeat breeding which extend their open period. These effects are attributed mainly to HS induced disturbances in luteinizing hormone (LH) secretion, follicular dominance and estrogen secretion, ovulation and oocyte competence, luteal development and progesterone secretion, utero-placental function and embryo-fetal development. Hormonal timed artificial insemination protocols and LH support around estrus improved summer pregnancy rates by avoiding need for estrus detection, assisting follicular development and ovulation, enhancing quality oocytes and stimulating luteal function. Progesterone supplementation to enhance embryonic development did not produce significant improvement in summer pregnancy rates. There is need for evaluating integrated approaches combining hormones, metabolic modifier and cyto-protective agents.

**Keywords:** Bovines, Heat stress, Hormones, Infertility.

### Introduction

Cows and buffaloes under summer heat stress (HS) face depressed reproductive activity. Summer infertility (SI) is characterized by anestrus, poor estrus expression and reduced fertility on breeding (Nardone *et al.*, 2010). SI is more important in tropical and sub-tropical areas, and in high yielding animals (Hansen, 2004). With current trends of rising global temperature, negative impact of SI on dairy production is expected to increase.

Current evidence indicates that HS and associated physiological changes hold immediate and delayed negative effects on: gonadotrophins secretion, follicular dynamics, ovulation, corpus luteum (CL) development, steroidogenesis, oocyte developmental competence, embryonic survival, utero-placental function, lactation and post-natal development (Wolfenson *et al.*, 1997; Bilby *et al.*, 2008; Hansen, 2009).

Knowledge that major SI mechanisms involve endocrine disturbance led to the development of hormone based interventions. Along with proper husbandry, hormonal supplementation and synchronization protocols demonstrate promise in enabling quick return to pregnancy (Hansen, 2012). Integration with supportive and protective interventions to address other physiological

disturbances and cell injury could further improve response to hormonal SI management strategies.

This review aimed to summarize the current knowledge on the mechanisms of bovine SI and examine their implication for successful hormonal intervention strategies.

### **Bovine Heat Stress**

The bovine Thermoneutral Zone (TNZ) is between 16°C to 25°C. Within this range, animals maintain a physiological body temperature of 38.4°C to 39.1°C (Yousef, 1985). Ambient temperatures above 20°C to 25°C enhance heat gain beyond that lost from the body inducing a state of HS with elevated internal body temperature (Yousef, 1985; Kennedy, 1999; Vale, 2007; Sunil Kumar *et al.*, 2011).

Animals try to restore thermal balance through mechanisms which reduce heat production - gain (reduced food intake, metabolism and activity; shade seeking; etc) and/or enhance heat loss (increasing water intake; bathing in ponds or mud; increased sweating respiration and salivation; redistributing blood flow towards peripheral integuments; etc). Cellular response to HS involves increased production of heat shock proteins (HSP) like HSP-70 and antioxidants to limit protein damage (Basiricò *et al.*, 2011). Physiological adjustments in HS pose negative consequences on energy availability, water and ionic

\*Corresponding Author: B.U. Wakayo. College of Veterinary Medicine-Jigjiga University, P.O.Box 1020, Jigjiga Ethiopian Somali Regional State, Ethiopia. Tel.: +251911731254; Fax: +251-025-7755947/76. Email: [fikeurga@gmail.com](mailto:fikeurga@gmail.com)

balance, milk production and fertility (Kennedy, 1999; Sunil Kumar *et al.*, 2011).

#### **Reproductive Impact of Summer Heat Stress**

The major impact of HS on reproduction involves delaying a return to gestation due to decreased submission rate and low conception / pregnancy rates (Nardone *et al.*, 2010). In heat stressed cows, duration and intensity of estrus decreases and incidence of anoestrous and silent ovulation increases (Hansen and Aréchiga, 1999; Lucy, 2002).

Cold season conception rates of 40% to 60% declines to 10% or 20% during hot summer months (Cavestany *et al.*, 1985). HS during gestation can compromise fetal development, lactation and neonatal development as well as postpartum ovarian activity (Bilby *et al.*, 2008). Buffaloes are more prone to HS and assume a seasonal calving avoiding summer breeding (Vale, 2007; Singh *et al.*, 2013).

#### **Mechanisms of Summer Infertility**

Summer heat stress disrupts reproduction in two general ways i.e. physiological adjustments such as redistribution of blood flow and energy deficits compromise reproductive activity indirectly and elevated internal body temperature directly disrupts endocrine mechanisms and cellular physiology (ovarian cells, germ cells, embryo or other) (Wolfenson *et al.*, 2000). Following sections attempt to summarize impacts of HS at various stages of the reproductive process, examine relationships between such effects and outline implications on key fertility indicators.

#### **Effect on gonadotropin secretion**

Both tonic and surge luteinizing hormone (LH) levels were depressed by HS in cows with low circulating estradiol concentrations (Gilad *et al.*, 1993). Poor LH surge interferes with maturation and ovulation of dominant follicles whereas low tonic LH levels hinder luteinization (luteal development) (Wolfenson *et al.*, 2000) and inhibit follicular growth and turnover in cyclic cows (Sirard, 2001).

Gonadotropin releasing hormone (GnRH) induced follicle stimulating hormone (FSH) secretion was similarly depressed during HS. However, tonic FSH secretion in HS is elevated due probably to reduced inhibition of negative feedback from small follicles (Wolfenson *et al.*, 1993; Khodaei-Motlagh *et al.*, 2011). FSH change related fertility effects are considered less relevant (Wolfenson *et al.*, 1993; Edwards and Hansen, 1996). Increased circulating prolactin level was reported during HS which could lead to acyclicity and infertility (Alamer, 2011; Singh *et al.*, 2013).

#### **Effect on ovarian activity**

HS reduced follicular dominance by inducing multiple large (> 10 mm) follicles as well as prolonged dominance of ovulatory follicles (Hansen, 2009).

Normal follicular selection and dominance could be disturbed by high tonic FSH availability. LH and negative energy balance (NEB) during HS can prevent maturation and ovulation of dominant follicles. Multiple dominant follicles can increase the rates of double ovulation and twin conceptions (Hansen, 2009). Prolonged follicular dominance disrupts normal oocyte maturation (Eg. premature meiosis) and reduces developmental competence (Mihm *et al.*, 1994).

Follicular estradiol secretion is depressed under HS primarily due to reduced theca cell androstenedione production associated with low 17 $\alpha$ -hydroxylase expression. Reduced granulosa cells aromatase activity and viability also contribute to poor estradiol secretion (Badinga *et al.*, 1993; Wolfenson *et al.*, 2000; Khodaei-Motlagh *et al.*, 2011). Low estrogen secretion disrupts expression of estrus, gonadotropin surge, ovulation, transport of gametes and fertilization (Wolfenson *et al.*, 2000).

In case dominant follicle/s ovulate, subsequent plasma progesterone concentrations are reduced during HS (Khodaei-Motlagh *et al.*, 2011). This could be attributed to small size of ovulatory follicle/s and low tonic LH stimulation of luteinization and steroidogenesis. Moreover, *in vitro* studies have indicated that HS compromised the viability of small luteal cells which represent a major source of progesterone (Wolfenson *et al.*, 2000). Low progesterone secretion limits endometrial function and embryo development (Wolfenson *et al.*, 2000; Khodaei-Motlagh *et al.*, 2011).

#### **Effect on oocyte competence**

Summer HS reduces oocyte developmental competence by affecting growth and maturation (Ju, 2005; Hansen, 2009). This could reflect a direct effect on the oocyte or an indirect consequence to supportive follicular cell changes. Direct effects on oocytes include: increased oxidative damage and apoptotic cell death (Paula-Lopes and Hansen, 2002), irreversible changes on cytoskeleton and meiotic spindle which interferes with cell division (Hansen, 2002; Ju, 2005), reduced mRNA and protein reserves for early embryonic development and altered membrane integrity which affects signal transduction and protein transport (Zeron *et al.*, 2001; Hansen, 2002).

On the other hand, tangible differences in important gene transcript levels (progesterone and prostaglandin receptors, transcription factors Egr-1 and DNA binding protein, etc) were detected between follicles with competent vs. incompetent oocytes (Robert *et al.*, 2001). Moreover, prolonged follicular dominance leads to premature meiosis and aged oocytes having poor developmental prospect (Hansen, 2002).

Reduced oocyte competence results in poor fertility rate after service. Stress capable of inducing oocyte

lesions in early stages of follicular growth could impair fertility even long after its removal meaning that depressed summer fertility persists even to autumn months (Wolfenson *et al.*, 2000; Hansen, 2002; Ju, 2005).

#### **Effect on embryo and fetus**

HS is a recognized cause of embryonic mortality. Embryonic loss in HS may result from poor oocyte quality. Moreover, peri-implantation embryos in early cleavage are highly sensitive to direct deleterious effects of HS (Ealy *et al.*, 1993; Bilby *et al.*, 2008; Hansen, 2009). HS causes embryonic death by interfering with protein synthesis (Edwards and Hansen, 1996), causing widespread apoptosis (Paula-Lopes and Hansen, 2002) and oxidative cell damage (Wolfenson *et al.*, 2000; Hansen, 2009) and reducing interferon-tau production for signaling pregnancy recognition (Wolfenson *et al.*, 1993; Bilby *et al.*, 2008).

Exposure of post-implantation embryos (early organogenesis) and fetus to HS leads to various teratologies (Wolfenson *et al.*, 2000; Hansen, 2009).

Genotype, stage of development and presence of cytoprotective molecules in female reproductive tract determine embryonic response to elevated temperature (Hansen, 2007). Embryonic susceptibility to HS declines in late cleavage (8 cells to morula stages). This was attributed to developmentally regulated protective mechanisms like: increasing expression of heat shock proteins (recently questioned by Sakatani *et al.* (2012)), declining oxidant production, increasing cytoplasmic antioxidant (glutathione) concentration and improving capacity for regulated apoptosis (Hansen, 2002; Paula-Lopes and Hansen, 2002; Ju, 2005; Hansen, 2009). The principle gave basis for successful application of embryo transfer advanced stage embryos to avoid pregnancy losses in summer (Hansen, 2012).

#### **Effect on uterus and placenta**

Rise in internal body temperature causes re-distribution of blood flow away from uterus which hampers supply of nutrients and hormones to the conceptus. Endometrial PGF-2 $\alpha$  secretion tends to increase in HS thereby threatening pregnancy maintenance (Vale, 2007; Bilby *et al.*, 2008). Placental weight and hormone secretion are reduced whereas vascular resistance is increased during HS. These effects further reduce perfusion of nutrients to the fetus and impair lactation (Hansen, 2009). Retarded fetal development due to inadequate nutrition can manifest as physiological and performance problems long after birth (Wolfenson *et al.*, 2000; Hansen, 2009).

#### **Hormonal Intervention to Summer Infertility**

Considering the role of endocrine disturbances in delaying service and hampering conception, hormonal

intervention presents a logical response for management of SI (Khodaei-Motlagh *et al.*, 2011). Multiple principles and protocols have been tried.

- Timed artificial insemination (TAI) protocols like OvSynch (GnRH – PGF-2 $\alpha$  - GnRH) help to circumvent estrus detection, resolve anestrus and silent cycle constraints, facilitate LH support to follicular and oocyte development. Such protocols helped to increase pregnancy rates and reduce open periods in HS. Replacing GnRH with human Chorionic Gonadotropin (hCG) (Hansen and Aréchiga, 1999) and simultaneous supplementation with cryoprotective agents (such as bovine somatotropin and antioxidants) further improved response to TAI strategies (Aréchiga *et al.*, 1998; Hansen, 2007). Dirandeh (2014) recently reported better outcome in heat stressed cows when ovsynch was initiated 6 days after estrus detected post the 30<sup>th</sup> day postpartum. The approach enhanced response to first GnRH, an important determinant of both synchronization and fertility on ovsynch TAI.

- Administration of GnRH during early estrus coinciding with endogenous LH surge enhances LH surge and improves synchrony between estrus-LH surge-ovulation-insemination. Timely ovulation, competent oocytes and enhanced luteal progesterone secretion during first 30 days of gestation improves fertilization and embryo survival. Better summer pregnancy rates of 18% to 29 % were reported using this approach (Wolfenson *et al.*, 2000).

- Progesterone supplementation at critical stages of embryonic development was another alternative for improving summer pregnancy rates. Induction of accessory corpora lutea after breeding (by ovulatory doses of hCG or GnRH) and long term application of progesterone supplements by controlled intravaginal drug releasing (CIDR) device to inseminated animals represent the main strategies evaluated. Despite their theoretical soundness, these strategies have thus far failed to unequivocally prove beneficial for improving summer pregnancy rates (Wolfenson *et al.*, 2000).

#### **Conclusion**

SI hinders profitable milk production and the problem is escalating with raising global temperatures. Mechanisms underlying SI involve disturbances in development and viability of follicular, luteal, germ, conceptus, endometrial and placental cells. These disturbances are driven by endocrine and energy deficits, physical, oxidative and apoptotic cell injuries and reduced protein synthesis and function. Developments on hormonal interventions assist to alleviate or circumvent certain aspects of SI. Further molecular insights on mechanisms underlying SI can lead to more novel management approaches. Lasting solution to SI requires creating more adaptable dairy herds and/or improving the production environment.

## References

- Alamer, M. 2011. The role of Prolactin in thermoregulation and water balance during heat stress in domestic animals. *Asian J. Anim. Vet. Adv.* (12), 1153-1169.
- Aréchiga, C.F., Staples, C.R., McDowell, L.R. and Hansen, P.J. 1998. Effects of timed insemination and supplemental betacarotene on reproduction and milk yield of dairy cows under heat stress. *J. Dairy Sci.* 81(2), 390-402.
- Badinga, L., Thatcher, W.W., Diaz, T., Drost, M. and Wolfenson, D. 1993. Effect of environmental heat stress on follicular development and steroidogenesis in lactating Holstein cows, *Theriogenology* 39, 797-810.
- Basiricò, L., Morera, P., Primi, V., Lacetera, N., Nardone, A. and Bernabucci, U. 2011. Cellular thermotolerance is associated with heat shock protein 70.1 genetic polymorphisms in Holstein lactating cows. *Cell Stress Chaperon* 16, 441-448.
- Bilby, T.R., Baumgard, L.H., Collier, R.J., Zimelman, R.B. and Rhoads, M.L. 2008. Heat stress effects on fertility: Consequences and possible solutions. In the Proceedings of the 2008 South Western Nutritional Conference.
- Cavestany, D., El-Whishy, A.B. and Foot, R.H. 1985. Effect of season and high environmental temperature on fertility of Holstein cattle. *J. Dairy Sci.* 68(6), 1471-1478.
- Dirandeh, E. 2014. Starting Ovsynch protocol on day 6 of first postpartum estrous cycle increased fertility in dairy cows by affecting ovarian response during heat stress. *Anim. Reprod. Sci.* 149(3-4), 135-140.
- Ealy, A.D., Drost, M. and Hansen, P.J. 1993. Developmental changes in embryonic resistance to adverse effects of maternal heat stress in cows. *J. Dairy Sci.* 76, 2899-2905.
- Edwards, J.L. and Hansen, P.J. 1996. Elevated temperature increases heat shock protein 70 synthesis in bovine two-cell embryos and compromises function of maturing oocytes. *Biol. Reprod.* 55, 340-346.
- Gilad, E., Meidan, R., Berman, A., Graber, Y. and Wolfenson, D. 1993. Effect of heat stress on tonic and GnRH-induced gonadotrophin secretion in relation to concentration of oestradiol in plasma of cyclic cows. *J. Reprod. Fertil.* 99, 315-321.
- Hansen, P.J. 2012. Solutions to Infertility Caused by Heat Stress. eXtension: <http://www.extension.org/pages/64288/solutions-to-infertility-caused-by-heat-stress>
- Hansen, P.J. 2009. Effects of heat stress on mammalian reproduction. *Phil. Trans. R. Soc. B.* 364, 3341-3350.
- Hansen, P.J. 2007. Exploitation of genetic and physiological determinants of embryonic resistance to elevated temperature to improve embryonic survival in dairy cattle during heat stress. *Theriogenology* 1(68), Suppl. 1, 242-249.
- Hansen, P.J. 2004. Physiological and cellular adaptations of zebu cattle to thermal stress. *Anim. Reprod. Sci.* 82-83, 349-360.
- Hansen, P.J. 2002. Embryonic mortality in cattle from the embryo's perspective. *J. Anim. Sci.* 80, 33-44.
- Hansen, P.J. and Aréchiga, C.F. 1999. Strategies for managing reproduction in the heat stressed. *J. Biol. Sci.* 1(1), 1-8.
- Ju, J.C. 2005. Cellular responses of oocytes and embryos under thermal stress: hints to molecular signaling. *Anim. Reprod.* 2(2), 79-90.
- Kennedy, B.S. 1999. Thermoregulation and the Effects of Heat Stress on Dairy Cattle. Mississippi State University College of Veterinary Medicine, Production Medicine Graduate Program. (<http://gpvec.unl.edu/heatdrought/HSDairyReview.htm>)
- Khodaei-Motlagh, M.M., Zare Shahneh, A., Masoumi, R. and Fabio, D. 2011. Alterations in reproductive hormones during heat stress in dairy cattle. *Afr. J. Biotechnol.* 10(29), 5552-5558.
- Lucy, M.C. 2002. Reproductive loss in farm animals during heat stress. In the Proceeding of 2002 15<sup>th</sup> Conference on Biometeorology and Aerobiology and the 16<sup>th</sup> International Congress of Biometeorology, pp: 50-53.
- Mihm, M., Baguisi, A., Boland, M.P. and Roche, J.F. 1994. Association between the duration of dominance of the ovulatory follicle and pregnancy rate in beef heifers. *J. Reprod. Fertil.* 102, 123-130.
- Nardone, A., Ronchi, B., Lacetera, N., Ranieri, M.S. and Bernabucci, U. 2010. Effects of climate changes on animal production and sustainability of livestock systems. *Livest. Sci.* 130, 57-69.
- Paula-Lopes, F.F. and Hansen, P.J. 2002. Apoptosis is an adaptive response in bovine preimplantation embryos that facilitates survival after heat shock. *Biochem. Biophys. Res. Co.* 295, 37-42.
- Robert, C., Gagne, D., Bousquet, D., Barnes, F.L. and Sirard, M.A. 2001. Differential display and suppressive subtractive hybridization used to identify granulosa cell messenger RNA associated with bovine oocyte developmental competence. *Biol. Reprod.* 64(6), 1812-1820.
- Sakatani, M., Alvarez, N.V., Takahashi, M. and Hansen, P.J. 2012. Consequences of physiological heat shock beginning at the zygote stage on embryonic development and expression of stress response genes in cattle. *J. Dairy Sci.*, 95(6), 3080-3091.

- Singh, M., Chaudhari, B.K., Singh, J.K., Singh, A.K. and Maurya, P.K. 2013. Effects of thermal load on buffalo reproductive performance during summer season. *J. Biol. Sci.* 1(1), 1-8.
- Sirard, M.A. 2001. Resumption of meiosis: mechanism involved in meiotic progression and its relation with developmental competence. *Theriogenology* 55, 1241-1254.
- Sunil Kumar, B.V., Kumar, A. and Kataria, M. 2011. Effect of heat stress in tropical livestock and different strategies for its amelioration. *J. Stress Physiol. Biochem.* 7 (1), 45-54.
- Vale, W.G. 2007. Effects of environment on buffalo reproduction. *Italian J. Anim. Sci.* 6(2), 130-142.
- Wolfenson, D., Bartol, F.F., Badinga, L., Barros, C.M., Marple, D.N., Cummings, K., Wolfe, D., Lucy, M.C., Spencer, T.E. and Thatcher, W.W. 1993. Secretion of PGF<sub>2a</sub> and oxytocin during hyperthermia in cyclic and pregnant heifers. *Theriogenology* 39, 1129-1141.
- Wolfenson, D., Lew, B.J., Thatcher, W.W., Graber, Y. and Meidan, R. 1997. Seasonal and acute heat stress effects on steroid production by dominant follicles in cow. *Anim. Reprod. Sci.* 47, 9-19.
- Wolfenson, D., Roth, Z. and Meidan, R. 2000. Impaired reproduction in heat stressed cattle: basic and applied aspects. *Anim. Reprod. Sci.* 60-61, 535-547.
- Yousef, M.K. 1985. *Stress Physiology in Livestock*. Vol.1. Boca Raton: CRC Press, pp: 67-73.
- Zeron, Y., Ocheretny, A., Kedar, O., Borochoy, A., Sklan, D. and Arav, A. 2001. Seasonal changes in bovine fertility: relation to developmental competence of oocytes, membrane properties and fatty acid composition of follicles. *Reprod.* 121, 447-454.